



Agromet Advisory Bulletin for Ernakulam District

(Valid from 03.02.2024 to 07.02.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 05/2024, (10) Friday

Dated 02.02.2024

A. Medium range weather forecast of IMD for next five days

Weather parameters	03.02.2024	04.02.2024	05.02.2024	06.02.2024	07.02.2024
Rainfall (mm)	0	0.2	0	0	0
Maximum temperature (°C)	32	32	32	32	32
Minimum temperature (°C)	25	26	26	26	26
Maximum Relative Humidity (%)	83	83	83	80	80
Minimum Relative Humidity (%)	51	51	51	50	50
Wind speed (kmph)	2	2	4	4	3
Wind direction (deg)	270	250	270	230	250
Total cloud cover (octa)	5	5	5	2	1

Weather summary / Alert: Since dry weather condition is prevailed, irrigation of crops can be continued, when required. Below normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

C. General and Agrometeorological Advisories for Ernakulum District (RRS Vyttila)

Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease	Crop/ Animal Advisory
Rice (Mundakan)	Harvesting time	Chaffy panicles to be removed from seed production plots. Field to be ploughed thoroughly after harvest. Preparations for cowpea or vegetable cultivation can be initiated in the fields where sufficient moisture is there.
Vegetables	Mites	2 per cent neem oil + garlic emulsion or commercial neem formulation azadirachtin 0.004 per cent (4 ml per litre of water) or micronized wettable sulphur 0.4 per cent can be applied in the initial stages of infestation. In the case of very severe infestation, the crop can be protected by spraying 0.05% dimethoate or Spiromesifen 22.9% SC 8 ml per 10 litre or Diafenthiuron 50% WP 1.2 g per 10 litre can be sprayed. Rational rotation of pesticides to be followed to avoid resistance development.
Vegetables (cowpea)	Fruit borer	To control fruit borer in cowpea, apply 5% Neem Kernel Suspension to control fruit borer in the initial stage. If the attack is severe, spray Flubendamide 2 ml per ten litre of water or Chlorantraniliprole 3 ml per ten litre of water.

Fisheries: - Farmers can culture the following fishes like red snapper, seabass, pearl spot, mullet, pompano, milk fish etc. in brackish water.