


Agromet Advisory Service bulletin 26/16, Tuesday dt 21-06-16

(Issued jointly by IAAS Vellanikkara)





ERNAKULAM

CURRENT SYNOPTIC SITUATION:

A trough of low pressure on sea level lies over East Arabian Sea off Karnataka-Kerala coast .An upper air cyclonic circulation lies over coastal Karnataka and neighbourhood between 3.6 and 5.8km a.s.l.

Heavy rainfall warning: Isolated Heavy rainfall is likely to occur over Kerala till the morning of 26th June.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 21.06.2016

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1	DAY-2	DAY-3	DAY-4	DAY-5
	22/06	23/06	24/06	25/06	26/06
ISTRICT : ERNAKULAM					
ainfall (mm)	20	20	30	20	 25
ax Temperature (deg C)	31	31	31	31	31
in Temperature (deg C)	24	24	24	24	24
otal cloud cover (octa)	6	7	8	6	5
ax Relative Humidity (%)	90	95	95	90	90
in Relative Humidity (%)	70	75	75	70	70
ind speed (kmph)	010	013	013	013	013
ind direction (deg)	290	290	290	290	290

Agro Meteorological Advisories:

Crop	Stage of the crop /	Advisories
	Pest/Disease	
Paddy (Jyothi, Kanchana)	Seedling stage	Due to high relative humidity, there is a chance of sheath blight, sheath rot, leaf spot diseases in paddy. This diseases can be control to an extend by dipping the roots of seedling in pseudomonas solution by mixing 20 g pseudomonas culture in one liter of water for 30 minutes while transplanting.
Coconut	Root wilt	Root wilt disease is seen in coconut. These coconut palms can be recovered by better management such as application of fertilizers, usage of high yielding resistant varieties instead of severely affected palms.
Pepper	Quick wilt	There is a chance of Quick wilt disease due to high relative humidity. Apply trichoderma or pseudomonas culture in soil or 1% Bordeaux mixture can be sprayed on the leaves and small branches.
Vegetables	White fly	Hang yellow boards dipped in caster oil in the field.