



Agromet Advisory Bulletin for PALAKKAD District

(Valid from 12.04.2025 to 16.04.2025)

*Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi*

AAS Bulletin No: 15/2025, (29) Friday

Dated 11.04.2025

A. Medium range weather forecast of IMD for next five days

Weather parameters	12.04.2025	13.04.2025	14.04.2025	15.04.2025	16.04.2025
Rainfall (mm)	1	0.4	8	7	3
Maximum temperature (°C)	37	37	37	37	37
Minimum temperature (°C)	26	26	26	26	26
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	55	55	55	55	55
Wind speed (kmph)	8	3	4	2	2
Wind direction (deg)	230	270	230	230	230
Total cloud cover (octa)	8	8	8	8	8

Weather summary / Alert: -. As per the forecast of IMD, there is a chance of light rainfall in the district.. Above normal rainfall is expected in Kerala.

Thunderstorm accompanied by lightning and gusty winds is very likely to occur at one or two places over Kerala on 11th to 15th April 2025.

B. Agrometeorological Advisories for crop protection & management

General and Agrometeorological Advisories for Palakkad District

General Advisory: - Proper mulching can be followed to conserve soil moisture. Since air temperature is increasing, it is a desirable practice to bury fresh or dried coconut husk in the coconut basin to conserve soil moisture

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Puncha rice (Dough stage)	Sheath Blight	To control sheath blight disease in rice, spray 4 gram of Trifloxystrobin + Tebuconazole mixture in ten litres of water.
Rice (virippu)	Management	It is recommended to sow dhaincha seeds in the paddy field before virippu planting.
Okra (Bhindi)	Mosaic disease	There is a chance of mosaic disease in bhindi. It can be controlled by removing the infected plants. Place yellow sticking traps in field to control the insect vector. If the attack is severe, spray Dimethoate 30 EC @ 1.5 ml per one litre of water.
Pepper	Slow wilt	To control the slow wilt disease in pepper, apply neem cake at the base of pepper plants. Also apply 25 grams of the bio control agent <i>Pisilomyces lilacinus</i> at the base of each plant. If the attack is severe, drench the basin of each plant with three gram Copper Oxy Chloride (COC) per one litre of water.

Animal Husbandry: - During cloudy condition, avoid bathing the cow and washing the cattle shed, otherwise more heat will be build up. Provide proper air circulation using fans in cattle shed. Assure increased ventilation in the cattle shed. Ensure sufficient supply of clean drinking water to the animals. (CAADECCS, KVASU, Mannuthy).