











Dated 11.02.2025

O (Dated 05t			uring last of 11 th Fe		025)		W 41 D 4				orecast Valid for 12 th to 16 th February 2025			
05/02	06/02	07/02	08/02	09/02	10/02	11/02	Date		12/02	13/02	14/02	15/02	16/02	
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)		0	0	0	0	0	
36.0	35.4	34.7	35.3	35.0	35.3	34.4	$T_{max}(^{0}C)$		34	34	34	34	34	
21.7	21.8	21.9	21.1	21.0	23.0	22.8	T mir	$n^{(0}C)$	23	23	23	23	23	
Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	AM	Cloud cover	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	
Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	PM							
73	71	78	64	73	69	67	RH-	[(%)	63	63	63	63	63	
27	33	23	26	29	34		RH-II (%)		22	22	22	22	22	
3.9	4.2	4.8	4.4	5.2	5.5	5.1	Wind Spe	ed (km/hr.)	4	2	2	2	2	
ESE	ESE	NNE	NNE	NNE	ENE	NNE	AM	Wind	W	CW	CVI	w	W	
ENE	NE	NNE	NNE	NNE			PM	direction	VV	SW	SW	VV	VV	
Rainfall (mm) in last week Rainfall (mm) from 0						/01/2025 to	till dated	Rainfall (mm) from 01/01/2025 to today						
0.0				0.0				0.0						

Weather summary / Alert: Since dry weather condition is prevails, irrigation of crops can be continued as per requirement. Below normal rainfall is expected in Kerala.

General and Agrometeorological Advisories for Thrissur District

General Advisory: - Prophylactic measures can be taken to control and to avoid the spread of crop pest and diseases. Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables. Proper mulching can be followed in order to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory				
Rice (Mundakan) Maturity stage	Rice bug	There is a chance of appearance of rice bug in paddy. To control spray fish jaggery extract (20 ml/litre of water) or spray chitin based Pseudomonas flurescens 20 gram in one litre of water. If the attack is severe, spray Azadirachtin 10 ml or Malathion 2 ml per litre of water.				
Brinjal	Shoot and Fruit Borer	There is a chance of shoot and fruit borer attack in brinjal. To control them, remove damaged shoot and fruits from the field. If the attack is severe, spray Chlorantraniliprole @ 3ml per ten litre of water.				
Chilli	Mite	There is a chance for the attack of mites in chilly. Apply Spiromesifen 8 ml per10 litre of water.				

Animal Husbandry: Ensure sufficient supply of clean drinking water to the animals. Allow the animal to graze early in the morning or in the evening to avoid direct sunlight. It is better to graze in the evening itself. (CAADECCS, KVASU, Mannuthy).